

LECTURE 18

Markov Processes – III

Readings: Section 7.4

Lecture outline

- Review of steady-state behavior
- Probability of blocked phone calls
- Calculating absorption probabilities
- Calculating expected time to absorption

Review

- Assume a single class of recurrent states, aperiodic; plus transient states. Then,

$$\lim_{n \rightarrow \infty} r_{ij}(n) = \pi_j$$

where π_j does not depend on the initial conditions:

$$\lim_{n \rightarrow \infty} \mathbf{P}(X_n = j \mid X_0 = i) = \pi_j$$


- π_1, \dots, π_m can be found as the unique solution to the balance equations

$$\pi_j = \sum_k \pi_k p_{kj}, \quad j = 1, \dots, m,$$

together with

$$\sum_j \pi_j = 1$$

Example




$$\pi_1 = 2/7, \pi_2 = 5/7$$

- Assume process starts at state 1.
- $\mathbf{P}(X_1 = 1, \text{ and } X_{100} = 1) =$
- $\mathbf{P}(X_{100} = 1 \text{ and } X_{101} = 2) =$

The phone company problem

- Calls originate as a Poisson process, rate λ
 - Each call duration is exponentially distributed (parameter μ)
 - B lines available
- Discrete time intervals of (small) length δ




- Balance equations: $\lambda \pi_{i-1} = i \mu \pi_i$

$$\pi_i = \pi_0 \frac{\lambda^i}{\mu^i i!} \quad \pi_0 = 1 / \sum_{i=0}^B \frac{\lambda^i}{\mu^i i!}$$

Calculating absorption probabilities

- What is the probability a_i that: process eventually settles in state 4, given that the initial state is i ?




For $i = 4$, $a_i =$

For $i = 5$, $a_i =$

$$a_i = \sum_j p_{ij} a_j, \quad \text{for all other } i$$

– unique solution

Expected time to absorption



- Find expected number of transitions μ_i , until reaching the absorbing state, given that the initial state is i ?

$\mu_i = 0$ for $i =$

For all other i : $\mu_i = 1 + \sum_j p_{ij} \mu_j$

– unique solution

Mean first passage and recurrence times

- Chain with one recurrent class; fix s recurrent

- **Mean first passage time from i to s :**

$$t_i = \mathbb{E}[\min\{n \geq 0 \text{ such that } X_n = s\} | X_0 = i]$$

- t_1, t_2, \dots, t_m are the unique solution to

$$t_s = 0, \\ t_i = 1 + \sum_j p_{ij} t_j, \quad \text{for all } i \neq s$$

- **Mean recurrence time of s :**

$$t_s^* = \mathbb{E}[\min\{n \geq 1 \text{ such that } X_n = s\} | X_0 = s]$$

- $t_s^* = 1 + \sum_j p_{sj} t_j$

MIT OpenCourseWare
<http://ocw.mit.edu>

6.041SC Probabilistic Systems Analysis and Applied Probability
Fall 2013

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.