

# CHAPTER 14 MULTIPLE INTEGRALS

## 14.1 Double Integrals (page 526)

The double integral  $\iint_R f(x, y) dA$  gives the volume between  $R$  and the surface  $z = f(x, y)$ . The base is first cut into small squares of area  $\Delta A$ . The volume above the  $i$ th piece is approximately  $f(x_i, y_i)\Delta A$ . The limit of the sum  $\sum f(x_i, y_i)\Delta A$  is the volume integral. Three properties of double integrals are  $\iint (f + g)dA = \iint f dA + \iint g dA$  and  $\iint c f dA = c \iint f dA$  and  $\iint_R f dA = \iint_S f dA + \iint_T f dA$  if  $R$  splits into  $S$  and  $T$ .

If  $R$  is the rectangle  $0 \leq x \leq 4, 4 \leq y \leq 6$ , the integral  $\iint x dA$  can be computed two ways. One is  $\iint x dy dx$ , when the inner integral is  $xy|_4^6 = 2x$ . The outer integral gives  $x^2|_0^4 = 16$ . When the  $x$  integral comes first it equals  $\int x dx = \frac{1}{2}x^2|_0^4 = 8$ . Then the  $y$  integral equals  $8y|_4^6 = 16$ . This is the volume between the base rectangle and the plane  $z = x$ .

The area  $R$  is  $\iint 1 dy dx$ . When  $R$  is the triangle between  $x = 0, y = 2x$ , and  $y = 1$ , the inner limits on  $y$  are  $2x$  and  $1$ . This is the length of a thin vertical strip. The (outer) limits on  $x$  are  $0$  and  $\frac{1}{2}$ . The area is  $\frac{1}{4}$ . In the opposite order, the (inner) limits on  $x$  are  $0$  and  $\frac{1}{2}y$ . Now the strip is horizontal and the outer integral is  $\int_0^1 \frac{1}{2}y dy = \frac{1}{4}$ . When the density is  $\rho(x, y)$ , the total mass in the region  $R$  is  $\iint \rho dx dy$ . The moments are  $M_y = \iint \rho x dx dy$  and  $M_x = \iint \rho y dx dy$ . The centroid has  $\bar{x} = M_y/M$ .

- $$1 \frac{8}{3}; \frac{2}{3} \quad 3 1; \ln \frac{3}{2} \quad 5 2 \quad 7 \frac{1}{2} \quad 9 \frac{4}{3} \quad 11 \int_{y=1}^2 \int_{x=1}^2 dx dy + \int_{y=2}^4 \int_{x=y/2}^2 dx dy$$
- 13**  $\int_{y=0}^1 \int_{x=-\frac{1}{2}\ln y}^{-\ln y} dx dy \quad 15 \int_{x=0}^1 \int_{y=-\sqrt{x}}^{\sqrt{x}} dy dx \quad 17 \int_0^1 \int_0^{y/2} dx dy = \int_0^{1/2} \int_{2x}^1 dy dx = \frac{1}{4}$
- 19**  $\int_0^3 \int_y^y dx dy = \int_{-1}^0 \int_{-x}^3 dy dx + \int_0^1 \int_x^3 dy dx = 9 \quad 21 \int_0^4 \int_{y/2}^y dx dy + \int_4^8 \int_{y/2}^4 dx dy = \int_0^4 \int_x^{2x} dy dx = 8$
- 23**  $\int_0^1 \int_0^{bx} dy dx + \int_1^2 \int_0^{b(2-x)} dy dx = \int_0^b \int_{y/b}^{2-(y/b)} dx dy = b \quad 25 f(a, b) - f(a, 0) - f(0, b) + f(0, 0)$
- 27**  $\int_0^1 \int_0^1 (2x - 3y + 1) dx dy = \frac{1}{2} \quad 29 \int_a^b f(x) dx = \int_a^b \int_0^{f(x)} 1 dy dx \quad 31 50,000\pi$
- 33**  $\int_1^3 \int_1^2 x^2 dx dy = \frac{14}{3} \quad 35 2 \int_0^{1/\sqrt{2}} \int_0^{\sqrt{1-y^2}} 1 dx dy = \frac{\pi}{4}$
- 37**  $\frac{1}{n^2} \sum_{j=1}^n \sum_{i=1}^n f\left(\frac{i-\frac{1}{2}}{n}, \frac{j-\frac{1}{2}}{n}\right)$  is exact for  $f = 1, x, y, xy \quad 39$  Volume 8.5  $\quad 41$  Volumes  $\ln 2, 2 \ln(1 + \sqrt{2})$
- 43**  $\int_0^1 \int_0^1 x^y dx dy = \int_0^1 \frac{1}{y+1} dy = \ln 2; \int_0^1 \int_0^1 x^y dy dx = \int_0^1 \frac{x-1}{\ln x} dx = \ln 2$
- 45** With long rectangles  $\sum y_i \Delta A = \sum \Delta A = 1$  but  $\iint y dA = \frac{1}{2}$

$$2 \int_1^e 2xy dx = x^2 y |_1^e = (e^2 - 1)y; \int_2^{2e} (e^2 - 1)y dy = (e^2 - 1) \frac{y^2}{2} |_2^{2e} = (e^2 - 1)(2e^2 - 2) = 2(e^2 - 1)^2;$$

$$\int_1^e \frac{dx}{xy} = \frac{\ln x}{y} |_1^e = \frac{1}{y}; \int_2^{2e} \frac{dy}{y} = \ln 2e - \ln 2 = \ln \frac{2e}{2} = 1.$$

**4**  $\int_1^2 ye^{xy} dx = e^{xy} |_1^2 = e^{2y} - e^y; \int_0^1 (e^{2y} - e^y) dy = [\frac{1}{2}e^{2y} - e^y]_0^1 = \frac{1}{2}e^2 - e + \frac{1}{2}; \int_0^3 \frac{dy}{\sqrt{3+2x+y}} = 2\sqrt{3+2x+y}]_0^3 = 2\sqrt{6+2x} - 2\sqrt{3+2x};$  the  $x$  integral is  $[\frac{2}{3}(6+2x)^{3/2} - \frac{2}{3}(3+2x)^{3/2}]_1^1 = \frac{2}{3}8^{3/2} - \frac{2}{3}5^{3/2} - \frac{2}{3}4^{3/2} + \frac{2}{3}.$

Note!  $3 + 2x + y$  is not zero in the region of integration.

**6** The region is above  $y = x^3$  and below  $y = x$  (from 0 to 1). Area =  $\int_0^1 (x - x^3) dx = [\frac{x^2}{2} - \frac{x^4}{4}]_0^1 = \frac{1}{4}.$

**8** The region is below the parabola  $y = 1 - x^2$  and above its mirror image  $y = x^2 - 1$ .

Area =  $\int_{-1}^1 (1 - x^2 - x^2 + 1) dx = [2x - \frac{2}{3}x^3]_{-1}^1 = \frac{8}{3}.$

- 10 The area is all below the axis  $y = 0$ , where horizontal strips cross from  $x = y$  to  $x = |y|$  (which is  $-y$ ). Note that the  $y$  integral stops at  $y = 0$ . Area =  $\int_{-1}^0 \int_y^{-y} dx dy = \int_{-1}^0 -2y dy = [-y^2]_{-1}^0 = 1$ .
- 12 The strips in Problem 6 from  $y = x^3$  up to  $x$  are changed to strips from  $x = y$  across to  $x = y^{1/3}$ . The outer integral on  $y$  is by chance also from 0 to 1. Area =  $\int_0^1 (y^{1/3} - y) dy = [\frac{3}{4}y^{4/3} - \frac{1}{2}y^2]_0^1 = \frac{1}{4}$ .
- 14 Between the upper parabola  $y = 1 - x^2$  in Problem 8 and the  $x$  axis, the strips now cross from the left side  $x = -\sqrt{1-y}$  to the right side  $x = +\sqrt{1-y}$ . This half of the area is  $\int_0^1 \int_{-\sqrt{1-y}}^{\sqrt{1-y}} dx dy = \int_0^1 2\sqrt{1-y} dy = -\frac{4}{3}(1-y)^{3/2}]_0^1 = \frac{4}{3}$ . The other half has strips from left side to right side of  $y = x^2 - 1$  or  $x = \pm\sqrt{1+y}$ . This area is  $\int_{-1}^0 \int_{-\sqrt{1+y}}^{\sqrt{1+y}} dx dy$  (also  $\frac{4}{3}$ ).
- 16 The triangle in Problem 10 had sides  $x = y$ ,  $x = -y$ , and  $y = -1$ . Now the strips are vertical. They go from  $y = -1$  up to  $y = x$  on the left side: area =  $\int_{-1}^0 \int_{-1}^x dy dx = \int_{-1}^0 (x+1) dx = \frac{1}{2}(x+1)^2]_{-1}^0 = \frac{1}{2}$ . The strips go from  $-1$  up to  $y = -x$  on the right side: area =  $\int_0^1 \int_{-1}^{-x} dy dx = \int_0^1 (-x+1) dx = \frac{1}{2}$ . Check:  $\frac{1}{2} + \frac{1}{2} = 1$ .
- 18 The triangle has corners at  $(0,0)$  and  $(-1,0)$  and  $(-1,-1)$ . Its area is  $\int_{-1}^0 \int_0^{-x} dy dx = \int_0^1 \int_{-1}^{-y} dx dy (= \frac{1}{2})$ .
- 20 The triangle has corners at  $(0,0)$  and  $(2,4)$  and  $(4,4)$ . Horizontal strips go from  $x = \frac{y}{2}$  to  $x = y$ : area =  $\int_0^4 \int_{y/2}^y dx dy = 4$ . Vertical strips are of two kinds: from  $y = x$  up to  $y = 2x$  or to  $y = 4$ . Area =  $\int_0^2 \int_x^{2x} dy dx + \int_2^4 \int_x^4 dy dx = 2 + 2 = 4$ .
- 22 (Hard Problem) The boundary lines are  $y = \frac{1}{2}x$  from  $(-2,-1)$  to  $(0,0)$ , and  $y = -2x$  from  $(0,0)$  to  $(1,-2)$ , and  $y = \frac{-x-5}{3}$  or  $x = -3y - 5$  from  $(-2,-1)$  to  $(1,-2)$ . (This is the hardest one: note first the slope  $-\frac{1}{3}$ .) Vertical strips go from the third line up to the first or second: area =  $\int_{-2}^0 \int_{(-x-5)/3}^{x/2} dy dx + \int_0^1 \int_{(-x-5)/3}^{-2x} dy dx = \frac{5}{3} + \frac{5}{6} = \frac{5}{2}$ . Horizontal strips cross from the first or third lines to the second: area =  $\int_{-2}^{-1} \int_{-3y-5}^{-y/2} dx dy + \int_{-1}^0 \int_{-3y-5}^{-y/2} dx dy = \frac{5}{4} + \frac{5}{4} = \frac{5}{2}$ .
- 24 The top of the triangle is  $(a,b)$ . From  $x = 0$  to  $a$  the vertical strips lead to  $\int_0^a \int_{dx/c}^{bx/a} dy dx = [\frac{bx^2}{2a} - \frac{dx^2}{2c}]_0^a = \frac{ba}{2} - \frac{da^2}{2c}$ . From  $x = a$  to  $c$  the strips go up to the third side:  $\int_a^c \int_{dx/c}^{b+(x-a)(d-b)/(c-a)} dy dx = [bx + \frac{(x-a)^2(d-b)}{2(c-a)} - \frac{dx^2}{2c}]_a^c = b(c-a) + \frac{(c-a)(d-b)}{2} - \frac{dc}{2} + \frac{da^2}{2c}$ . The sum is  $\frac{ba}{2} + \frac{b(c-a)}{2} + \frac{d(c-a)}{2} - \frac{dc}{2} = \frac{bc-ad}{2}$ . This is half of a parallelogram.
- 26  $\int_0^b \int_0^a \frac{\partial f}{\partial x} dx dy = \int_0^b [f(a,y) - f(0,y)] dy$ .
- 28 Over the square  $\int_0^1 \int_0^1 (xe^y - ye^x) dy dx = \int_0^1 (xe - \frac{e^x}{2} - x) dx = [\frac{x^2e}{2} - \frac{e^x}{2} - \frac{x^2}{2}]_0^1 = \frac{e}{2} - \frac{e}{2} - \frac{1}{2} + \frac{1}{2} = 0$ . (Looking back: zero is not a surprise because of symmetry.) Over the triangle the integral up to  $y = x$  is  $\int_0^1 \int_0^x (xe^y - ye^x) dy dx$ . Over the triangle across to  $y = x$  the integral is  $\int_0^1 \int_0^y (xe^y - ye^x) dx dy$ . Exchange  $y$  and  $x$  in the second double integral to get *minus* the first double integral.
- 30  $\int_{-1}^1 (1-x^2) dx = [x - \frac{x^3}{3}]_{-1}^1 = \frac{4}{3}$ . With horizontal strips this is  $\int_0^1 \int_{-\sqrt{1-y}}^{\sqrt{1-y}} dx dy = \int_0^1 2\sqrt{1-y} dy = -\frac{4}{3}(1-y)^{3/2}]_0^1 = \frac{4}{3}$ .
- 32 The height is  $z = \frac{1-ax-by}{c}$ . Integrate over the triangular base ( $z = 0$  gives the side  $ax + by = 1$ ): volume =  $\int_{z=0}^{1/a} \int_{y=0}^{(1-az)/b} \int_{x=0}^{1-az-by} \frac{1-az-by}{c} dy dx = \int_0^{1/a} \frac{1}{c} [y - axy - \frac{1}{2}by^2]_0^{(1-az)/b} dx = \int_0^{1/a} \frac{1}{c} \frac{(1-az)^2}{2b} dx = -\frac{(1-az)^3}{6abc}]_0^{1/a} = \frac{1}{6abc}$ .
- 34 From Problem 33 the mass is  $\frac{14}{3}$ . The moments are  $\int_1^3 \int_1^2 x^3 dx dy = \int_1^3 \frac{2^4-1^4}{4} dy = \frac{15}{2}$  and  $\int_1^3 \int_1^2 yx^2 dx dy = \int_1^3 \frac{8-1}{3} y dy = \frac{28}{3}$ . Then  $\bar{x} = \frac{15/2}{14/3} = \frac{45}{28}$  and  $\bar{y} = \frac{28/3}{14/3} = 2$ .
- 36 The area of the quarter-circle is  $\frac{\pi}{4}$ . The moment is zero around the axis  $y = 0$  (by symmetry):  $\bar{x} = 0$ . The other moment, with a factor 2 that accounts for symmetry of left and right, is  $2 \int_0^{\sqrt{2}/2} \int_x^{\sqrt{1-x^2}} y dy dx = 2 \int_0^1 (\frac{1-x^2}{2} - \frac{x^2}{2}) dx = 2[\frac{x}{2} - \frac{x^3}{3}]_0^1 = \frac{\sqrt{2}}{3}$ . Then  $\bar{y} = \frac{\sqrt{2}/3}{\pi/4} = \frac{4\sqrt{2}}{3\pi}$ .
- 38 The integral  $\int_0^1 \int_0^1 x^2 dx dy$  has the usual midpoint error  $-\frac{(\Delta x)^2}{12}$  for the integral of  $x^2$  (see Section 5.8). The  $y$  integral  $\int_0^1 dy = 1$  is done exactly. So the error is  $-\frac{1}{12n^2}$  (and the same for  $\iint y^2 dx dy$ ). The integral of  $xy$  is computed exactly. Errors decrease with exponent  $p = 2$ , the order of accuracy.

40 The exact integral is  $\int_0^1 \int_0^1 \frac{dx dy}{\sqrt{x^2+y^2}} = 2 \int_0^{\pi/4} \int_0^{\sec \theta} \frac{r dr d\theta}{r} = 2 \int_0^{\pi/4} \sec \theta d\theta = 2 \ln(\sec \theta + \tan \theta) \Big|_0^{\pi/4} = 2 \ln(\sqrt{2} + 1)$ .

42 The exact integral is  $\int_0^1 \int_0^1 e^x \sin \pi y dx dy = \int_0^1 (e - 1) \sin \pi y dy = \frac{e-1}{\pi} (-\cos \pi y) \Big|_0^1 = \frac{2}{\pi}(e - 1)$ .

## 14.2 Change to Better Coordinates (page 534)

We change variables to improve the limits of integration. The disk  $x^2 + y^2 \leq 9$  becomes the rectangle  $0 \leq r \leq 3, 0 \leq \theta \leq 2\pi$ . The inner limits of  $\iint dy dx$  are  $y = \pm\sqrt{9 - x^2}$ . In polar coordinates this area integral becomes  $\iint r dr d\theta = 9\pi$ .

A polar rectangle has sides  $dr$  and  $r d\theta$ . Two sides are not straight but the angles are still  $90^\circ$ . The area between the circles  $r = 1$  and  $r = 3$  and the rays  $\theta = 0$  and  $\theta = \pi/4$  is  $\frac{1}{8}(3^2 - 1^2) = 1$ . The integral  $\iint x dy dx$  changes to  $\iint r^2 \cos \theta dr d\theta$ . This is the moment around the y axis. Then  $\bar{x}$  is the ratio  $M_y/M$ . This is the x coordinate of the centroid, and it is the average value of  $x$ .

In a rotation through  $\alpha$ , the point that reaches  $(u, v)$  starts at  $x = u \cos \alpha - v \sin \alpha, y = u \sin \alpha + v \cos \alpha$ . A rectangle in the  $uv$  plane comes from a rectangle in  $xy$ . The areas are equal so the stretching factor is  $J = 1$ . This is the determinant of the matrix  $\begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$ . The moment of inertia  $\iint x^2 dx dy$  changes to  $\iint (u \cos \alpha - v \sin \alpha)^2 du dv$ .

For single integrals  $dx$  changes to  $(dx/du)du$ . For double integrals  $dx dy$  changes to  $J du dv$  with  $J = \partial(x, y)/\partial(u, v)$ . The stretching factor  $J$  is the determinant of the 2 by 2 matrix  $\begin{bmatrix} \partial x/\partial u & \partial x/\partial v \\ \partial y/\partial u & \partial y/\partial v \end{bmatrix}$ . The functions  $x(u, v)$  and  $y(u, v)$  connect an  $xy$  region  $R$  to a  $uv$  region  $S$ , and  $\iint_R dx dy = \iint_S J du dv = \text{area of } R$ . For polar coordinates  $x = u \cos v$  and  $y = u \sin v$  (or  $r \sin \theta$ ). For  $x = u, y = u + 4v$  the 2 by 2 determinant is  $J = 4$ . A square in the  $uv$  plane comes from a parallelogram in  $xy$ . In the opposite direction the change has  $u = x$  and  $v = \frac{1}{4}(y - x)$  and a new  $J = \frac{1}{4}$ . This  $J$  is constant because this change of variables is linear.

$$1 \int_{\pi/4}^{3\pi/4} \int_0^1 r dr d\theta = \frac{\pi}{4} \quad 3 S = \text{quarter-circle with } u \geq 0 \text{ and } v \geq 0; \int_0^1 \int_0^{\sqrt{1-v^2}} du dv$$

$$5 R \text{ is symmetric across the } y \text{ axis; } \int_0^1 \int_0^{\sqrt{1-v^2}} u du dv = \frac{1}{3} \text{ divided by area gives } (\bar{u}, \bar{v}) = (4/3\pi, 4/3\pi)$$

$$7 2 \int_0^{1/\sqrt{2}} \int_{1+x}^{1+\sqrt{1-x^2}} dy dx; xy \text{ region } R^* \text{ becomes } R \text{ in the } x^*y^* \text{ plane; } dx dy = dx^*dy^* \text{ when region moves}$$

$$9 J = \begin{vmatrix} \partial x/\partial r^* & \partial x/\partial \theta^* \\ \partial y/\partial r^* & \partial y/\partial \theta^* \end{vmatrix} = \begin{vmatrix} \cos \theta^* & -r^* \sin \theta^* \\ \sin \theta^* & r^* \cos \theta^* \end{vmatrix} = r^*; \int_{\pi/4}^{3\pi/4} \int_0^1 r^* dr^* d\theta^*$$

$$11 I_y = \iint_R x^2 dx dy = \int_{\pi/4}^{3\pi/4} \int_0^1 r^2 \cos^2 \theta r dr d\theta = \frac{\pi}{16} - \frac{1}{8}; I_x = \frac{\pi}{16} + \frac{1}{8}; I_0 = \frac{\pi}{8}$$

$$13 (0,0), (1,2), (1,3), (0,1); \text{area of parallelogram is 1}$$

$$15 x = u, y = u + 3v + uv; \text{then } (u, v) = (1, 0), (1, 1), (0, 1) \text{ give corners } (x, y) = (1, 0), (1, 5), (0, 3)$$

$$17 \text{ Corners } (0,0), (2,1), (3,3), (1,2); \text{sides } y = \frac{1}{2}x, y = 2x - 3, y = \frac{1}{2}x + \frac{3}{2}, y = 2x$$

$$19 \text{ Corners } (1,1), (e^2, e), (e^3, e^3), (e, e^2); \text{sides } x = y^2, y = x^2/e^3, x = y^2/e^3, y = x^2$$

$$21 \text{ Corners } (0,0), (1,0), (1,2), (0,1); \text{sides } y = 0, x = 1, y = 1 + x^2, x = 0$$

23  $J = \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} = 3$ , area  $\int_0^1 \int_0^1 3du dv = 3$ ;  $J = \begin{vmatrix} 2e^{2u+v} & e^{2u+v} \\ e^{u+2v} & 2e^{u+2v} \end{vmatrix} = 3e^{3u+3v}, \int_0^1 \int_0^1 3e^{3u+3v} du dv = \int_0^1 (e^{3+3v} - e^{3v}) dv = \frac{1}{3}(e^6 - 2e^3 + 1)$

25 Corners  $(x, y) = (0, 0), (1, 0), (1, f(1)), (0, f(0)); (\frac{1}{2}, 1)$  gives  $x = \frac{1}{2}, y = f(\frac{1}{2}); J = \begin{vmatrix} 1 & 0 \\ vf'(u) & f(u) \end{vmatrix} = f(u)$

27  $B^2 = 2 \int_0^{\pi/4} \int_0^{1/\sin\theta} e^{-r^2} r dr d\theta = \int_0^{\pi/4} (e^{-1/\sin^2\theta} - 1) d\theta$

29  $F = \iint r^2 dr d\theta / \iint r dr d\theta = \int_0^{\pi} \frac{8}{3} a^3 \sin^3 \theta d\theta / \pi a^2 = \frac{32a}{9\pi}$       31  $\int_0^{2\pi} \int_0^1 r^2 r dr d\theta = \frac{\pi}{2}$

33 Along the right side; along the bottom; at the bottom right corner

35  $\iint xy dx dy = \int_0^1 \int_0^1 (u \cos \alpha - v \sin \alpha)(u \sin \alpha + v \cos \alpha) du dv = \frac{1}{4}(\cos^2 \alpha - \sin^2 \alpha)$

37  $\int_0^{2\pi} \int_4^5 r^2 r dr d\theta = \frac{2\pi}{6}(5^6 - 4^6)$       39  $x = \cos \alpha - \sin \alpha, y = \sin \alpha + \cos \alpha$  goes to  $u = 1, v = 1$

2 Area  $= \int_{-\sqrt{2}/2}^{\sqrt{2}/2} \int_{|x|}^{\sqrt{1-x^2}} dy dx$  splits into two equal parts left and right of  $x = 0 : 2 \int_0^{\sqrt{2}/2} \int_x^{\sqrt{1-x^2}} dy dx = 2 \int_0^{\sqrt{2}/2} (\sqrt{1-x^2} - x) dx = [x\sqrt{1-x^2} + \sin^{-1} x - x^2]_0^{\sqrt{2}/2} = \sin^{-1} \frac{\sqrt{2}}{2} = \frac{\pi}{4}$ . The limits on

$\iint dx dy$  are  $\int_0^{\sqrt{2}/2} \int_{-y}^y dx dy$  for the lower triangle plus  $\int_{\sqrt{2}/2}^1 \int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} dx dy$  for the circular top.

4 (See Problem 36 of Section 14.1)  $\int_{\pi/4}^{3\pi/4} \int_0^1 (r \sin \theta) r dr d\theta = [\frac{r^2}{3}]_0^1 [-\cos \theta]_{\pi/4}^{3\pi/4} = \frac{\sqrt{2}}{3}$ ; divide by area  $\frac{\pi}{4}$  to reach  $\bar{y} = \frac{\sqrt{2}/3}{\pi/4} = \frac{4\sqrt{2}}{3\pi}$ .

6 Area of wedge  $= \frac{b}{2\pi}(\pi a^2)$ . Divide  $\int_0^b \int_0^a (r \cos \theta) r dr d\theta = \frac{a^3}{3} \sin b$  by this area  $\frac{ba^2}{2}$  to find

$\bar{x} = \frac{2a}{3b} \sin b$ . (Interesting limit:  $\bar{x} \rightarrow \frac{2}{3}a$  as the wedge angle  $b$  approaches zero: like the centroid of a triangle.)

For  $\bar{y}$  divide  $\int_0^b \int_0^a (r \sin \theta) r dr d\theta = \frac{a^3}{3}(1 - \cos b)$  by the area  $\frac{ba^2}{2}$  to find  $\bar{y} = \frac{2a}{3b}(1 - \cos b)$ .

8 The limits on  $r, \theta$  are extremely awkward for  $R^*$ . Contrast with the simple limits  $0 \leq r^* \leq 1, \frac{\pi}{4} \leq \theta^* \leq \frac{3\pi}{4}$  when the coordinates are recentered at  $(0, 1)$ . (A point on the lower boundary of the wedge has  $r = \frac{\sin \frac{3\pi}{4}}{\sin(\frac{\pi}{4} - \theta)}$  by the law of sines.)

10 The centroid  $(0, \bar{y})$  of  $R$  moves up to the centroid  $(0, \bar{y} + 1)$  of  $R^*$ . The centroid of a circle is its center  $(1, 2)$ . The centroid of the upper half is  $(1, 2 + \frac{4}{\pi})$  because a half-circle has  $\int_0^{\pi} \int_0^3 (r \sin \theta) r dr d\theta = 18$  divided by its area  $\frac{9\pi}{2}$  (which gives  $\frac{4}{\pi}$ ).

12  $I_x = \int_{\pi/4}^{3\pi/4} \int_0^1 (r \sin \theta + 1)^2 r dr d\theta = \frac{1}{4} \int \sin^2 \theta d\theta + \frac{2}{3} \int \sin \theta d\theta + \frac{1}{2} \int d\theta = [\frac{\theta}{8} - \frac{\sin 2\theta}{16} - \frac{2}{3} \cos \theta + \frac{\theta}{2}]_{\pi/4}^{3\pi/4} = \frac{5\pi}{16} + \frac{2}{16} + \frac{4}{3} \frac{\sqrt{2}}{2}; I_y = \iint (r \cos \theta)^2 r dr d\theta = \frac{\pi}{16} - \frac{1}{8}$  (as in Problem 11);  $I_0 = I_x + I_y = \frac{3\pi}{8} + \frac{4}{3} \frac{\sqrt{2}}{2}$ .

14 The corner  $(1, 2)$  should be (a, c), when  $u = 0$  and  $v = 1$ ; the corner  $(0, 1)$  should be (b, d), when  $u = 1$  and  $v = 0$ . Check at  $u = v = 1$ ; there  $x = au + bv = 1$  and  $y = cu + dv = 3$  to give the correct corner  $(1, 3)$ .

Then  $J = ad - bc = (1)(1) - (0)(2) = 1$ . The unit square has area 1; so does  $R$ .

16 A linear change takes the square  $S$  into a parallelogram  $R$  (with one corner at  $(0, 0)$ ). Reason: The vector sum of the two sides from  $(0, 0)$  is still the vector to the far corner.

18 Corners when  $u = 0$  or 1,  $v = 0$  or 1:  $(0, 0), (3, 1), (5, 2), (2, 1)$ . The sides have equations

$$y = \frac{1}{3}x, y = \frac{1}{2}x - \frac{1}{2}, y = \frac{1}{3}x + \frac{1}{3}, y = \frac{1}{2}x.$$

20 Corners when  $u = 0$  or 1,  $v = 0$  or 1:  $(0, 0), (0, -1), (1, 0), (0, 1)$ . Actually  $(0, 0)$  is not a corner because one side comes down the  $y$  axis. The side with  $u = 1$  is  $x = v, y = v^2 - 1$  or  $y = x^2 - 1$ . The side with  $v = 1$  is  $x = u, y = 1 - u^2$  or  $y = 1 - x^2$ .

22 Here  $u = 0$  or 1,  $v = 0$  or 1 gives the corners  $(0, 0), (1, 0), (\cos 1, \sin 1)$ . The side with  $u = 1$  is a circular arc  $x = \cos v, y = \sin v$  between the last two corners. The other sides are straight: the region is pie-shaped (a fraction  $\frac{1}{2\pi}$  of the unit circle).

**24** Problem 18 has  $J = \begin{vmatrix} 3 & 2 \\ 1 & 1 \end{vmatrix} = 1$ . So the area of  $R$  is  $1 \times$  area of unit square = 1. Problem 20 has

$J = \begin{vmatrix} v & u \\ -2u & 2v \end{vmatrix} = 2(u^2 + v^2)$ , and integration over the square gives area of  $R$  =

$\int_0^1 \int_0^1 2(u^2 + v^2) du dv = \frac{4}{3}$ . Check in  $x, y$  coordinates: area of  $R = 2 \int_0^1 (1 - x^2) dx = \frac{4}{3}$ .

**26**  $\left| \begin{matrix} \partial r / \partial x & \partial r / \partial y \\ \partial \theta / \partial x & \partial \theta / \partial y \end{matrix} \right| = \left| \begin{matrix} x/r & y/r \\ -y/r^2 & x/r^2 \end{matrix} \right| = \frac{x^2 + y^2}{r^3} = \frac{1}{r}$ . As in equation 12, this new  $J$  is  $\frac{1}{\text{old } J}$ .

**28**  $\int_{-\infty}^{\infty} x^2 e^{-x^2/2} dx = (u)(v) - \int v du = (x)(-e^{-x^2/2}) \Big|_{-\infty}^{\infty} + \int_{-\infty}^{\infty} e^{-x^2/2} dx = 0 + \sqrt{2\pi}$  by Example 5. Divide by  $\sqrt{2\pi}$  to find  $\sigma^2 = 1$ .

**30**  $R$  is an infinite strip above the interval  $[0,1]$  on the  $x$  axis. Its boundary  $x = 1$  is  $r \cos \theta = 1$  or  $r = \sec \theta$ .

The limits are  $0 \leq r \leq \sec \theta$  and  $0 \leq \theta \leq \frac{\pi}{2}$ . The integral is  $\int_0^{\pi/2} \int_0^{\sec \theta} \frac{r dr d\theta}{r^3} = \int_0^{\pi/2} (\infty) d\theta = \infty$ .

For a finite example integrate  $(x^2 + y^2)^{-1/2} = \frac{1}{r}$ .

**32** Equation (3) with  $y$  instead of  $x$  has  $\iint y^2 dA = \int_0^1 \int_0^1 (u \sin \alpha + v \cos \alpha)^2 du dv = \sin^2 \alpha \iint u^2 du dv + \sin \alpha \cos \alpha \iint 2uv du dv + \cos^2 \alpha \iint v^2 du dv = \frac{\sin^2 \alpha}{3} + \frac{\sin \alpha \cos \alpha}{2} + \frac{\cos^2 \alpha}{3}$ .

**34** (a) False (forgot the stretching factor  $J$ ) (b) False ( $x$  can be larger than  $x^2$ ) (c) False (forgot to divide by the area) (d) True (odd function integrated over symmetric interval) (e) False (the straight-sided region is a trapezoid: angle from 0 to  $\theta$  and radius from  $r_1$  to  $r_2$  yields area  $\frac{1}{2}(r_2^2 - r_1^2) \sin \theta \cos \theta$ ).

**36**  $\iint \rho dA = \int_0^{2\pi} \int_4^5 r^2 (r dr d\theta) = 2\pi \frac{5^4 - 4^4}{4}$ . This is the polar moment of inertia  $I_0$  with density  $\rho = 1$ .

**38**  $\iint f dA = f(P) \iint dA$  is the Mean Value Theorem for double integrals (compare Property 7, Section 5.6). If  $f = x$  or  $f = y$ , choose  $P = \text{centroid } (\bar{x}, \bar{y})$ .

## 14.3 Triple Integrals (page 540)

Six important solid shapes are a **box**, **prism**, **cone**, **cylinder**, **tetrahedron**, and **sphere**. The integral  $\iiint dz dy dz$  adds the volume  $dx dy dz$  of small boxes. For computation it becomes three single integrals. The inner integral  $\int dz$  is the length of a line through the solid. The variables  $y$  and  $z$  are held constant. The double integral  $\iint dx dy$  is the area of a slice, with  $z$  held constant. Then the  $z$  integral adds up the volumes of slices.

If the solid region  $V$  is bounded by the planes  $x = 0, y = 0, z = 0$ , and  $x + 2y + 3z = 1$ , the limits on the inner  $x$  integral are 0 and  $1 - 2y - 3z$ . The limits on  $y$  are 0 and  $\frac{1}{2}(1 - 3z)$ . The limits on  $z$  are 0 and  $\frac{1}{3}$ . In the new variables  $u = x, v = 2y, w = 3z$ , the equation of the outer boundary is  $u + v + w = 1$ . The volume of the tetrahedron in  $uvw$  space is  $\frac{1}{6}$ . From  $dx = du$  and  $dy = dv/2$  and  $dz = dw/3$ , the volume of an  $xyz$  box is  $dx dy dz = \frac{1}{6} du dv dw$ . So the volume of  $V$  is  $\frac{1}{36}$ .

To find the average height  $\bar{z}$  in  $V$  we compute  $\iiint z dV / \iiint dV$ . To find the total mass if the density is  $\rho = e^z$  we compute the integral  $\iiint e^z dx dy dz$ . To find the average density we compute  $\iiint e^z dV / \iiint dV$ . In the order  $\iiint dz dx dy$  the limits on the inner integral can depend on  $x$  and  $y$ . The limits on the middle integral can depend on  $y$ . The outer limits for the ellipsoid  $x^2 + 2y^2 + 3z^2 \leq 8$  are  $-2 \leq y \leq 2$ .

$$1 \int_0^1 \int_0^z \int_0^y dx dy dz = \frac{1}{6}$$

3  $0 \leq y \leq x \leq z \leq 1$  and all other orders  $zxy, yzx, zxy, zyx$ ; all six contain  $(0,0,0)$ ; to contain  $(1,0,1)$

- 5  $\int_{-1}^1 \int_{-1}^1 \int_{-1}^1 dx dy dz = 8$       7  $\int_{-1}^1 \int_{-1}^1 \int_{-1}^1 dx dy dz = 4$       9  $\int_{-1}^1 \int_1^1 \int_1^z dx dy dz = \frac{4}{3}$   
 11  $\int_0^1 \int_0^{2-z} \int_0^{2-y-2z} dx dy dz = \frac{2}{3}$       13  $\int_0^{1/3} \int_0^{2-z} \int_0^{2-y-2z} dx dy dz = \frac{7}{12}$   
 15  $\int_0^1 \int_0^{1-z} \int_0^{\sqrt{(1-z)^2-y^2}} dx dy dz = \frac{\pi}{3}$       17  $\int_0^6 \int_0^1 \int_0^{\sqrt{1-y^2}} dx dy dz = 6\pi$       19  $\int_0^1 \int_0^1 \int_0^{\sqrt{1-y^2}} dx dy dz = \pi$   
 21 Corner of cube at  $(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}})$ ; sides  $\frac{2}{\sqrt{3}}$ ; area  $\frac{8}{3\sqrt{3}}$   
 23 Horizontal slices are circles of area  $\pi r^2 = \pi(4-z)$ ; volume  $= \int_0^4 \pi(4-z)dz = 8\pi$ ; centroid  
 has  $\bar{x} = 0, \bar{y} = 0, \bar{z} = \int_0^4 z\pi(4-z)dz/8\pi = \frac{4}{3}$   
 25  $I = \frac{z^2}{2}$  gives zeros;  $\frac{\partial I}{\partial x} = \int_0^x \int_0^y f dy dz, \frac{\partial I}{\partial y} = \int_0^x \int_0^x f dx dz, \frac{\partial^2 I}{\partial y \partial z} = \int_0^x f dx$   
 27  $\int_{-1}^1 \int_{-1}^1 \int_{-1}^1 (y^2 + z^2) dx dy dz = \frac{16}{3}; \iiint x^2 dV = \frac{8}{3}; 3 \iiint (x - \frac{x+y+z}{3})^2 dV = \frac{16}{3}$   
 29  $\int_0^3 \int_0^2 \int_0^y dx dy dz = 6$       31 Trapezoidal rule is second-order; correct for 1,  $x, y, z, xy, xz, yz, xyz$

- 2 The area of  $0 \leq x \leq y \leq z \leq 1$  is  $\int_0^1 \int_x^1 \int_y^1 dz dy dx$ . The four faces are  $x = 0, y = x, z = y, z = 1$ .  
 4  $\int_0^1 \int_0^x \int_0^y x dx dy dz = \int_0^1 \int_0^x \frac{y^2}{2} dy dz = \int_0^1 \frac{x^3}{6} dz = \frac{1}{24}$ . Divide by the volume  $\frac{1}{6}$  to find  $\bar{x} = \frac{1}{4}$ ;  
 $\int_0^1 \int_0^x \int_0^y y dx dy dz = \int_0^1 \int_0^x y^2 dy dz = \int_0^1 \frac{x^3}{3} dz = \frac{1}{12}$  and  $\bar{y} = \frac{1}{2}$ ; by symmetry  $\bar{z} = \frac{3}{4}$ .  
 6 Volume of half-cube  $= \int_0^1 \int_{-1}^1 \int_{-1}^1 dx dy dz = 4$ .  
 8  $\int_0^1 \int_{-1}^1 \int_{-1}^1 dx dy dz = \int_0^1 2(z+1)dz = [(z+1)^2]_0^1 = 3$ .  
 10  $\int_{-1}^1 \int_{-1}^1 \int_{-1}^y dx dy dz = \int_{-1}^1 \int_{-1}^y (y+1)dy dz = \int_{-1}^1 \frac{(z+1)^2}{2} dz = [\frac{(z+1)^3}{6}]_{-1}^1 = \frac{4}{3}$  (tetrahedron).  
 12 The plane faces are  $x = 0, y = 0, z = 0$ , and  $2x + y + z = 4$  (which goes through 3 points). The volume  
 is  $\int_0^2 \int_0^{4-2x} \int_0^{4-2x-y} dz dy dx = \int_0^2 \int_0^{4-2x} (4-2x-y) dy dx = \int_0^2 \frac{(4-2x)^2}{2} dx = [-\frac{(4-2x)^3}{12}]_0^2 = \frac{4^3}{12} = \frac{16}{3}$ .  
 Check: Multiply standard volume  $\frac{1}{6}$  by  $(4)(4)(2) = \frac{16}{3}$ . Check: Double the volume in Problem 11.  
 14 Put  $dz$  last and stop at  $z = 1$ :  $\int_0^1 \int_0^x \int_0^{(4-y-z)/2} dx dy dz = \int_0^1 \int_0^x \frac{4-y-z}{2} dy dz =$   
 $\int_0^1 \frac{(4-z)^2}{4} dz = [-\frac{(4-z)^3}{12}]_0^1 = \frac{4^3-3^3}{12} = \frac{37}{12}$ .  
 16 (Still tetrahedron of Problem 12: volume still  $\frac{16}{3}$ ). Limits of integration: the top vertex  
 falls from  $(0,0,4)$  onto the  $y$  axis at  $(0, -4, 0)$ . The corner  $(2,0,0)$  stays on the  $x$  axis.  
 The corner  $(0,4,0)$  swings up to  $(0,0,4)$ . The volume integral is  $\int_0^4 \int_{-4}^0 \int_0^x dx dy dz = \frac{16}{3}$ .  
 18 The plane  $z = x$  cuts the circular base in half, leaving  $x \geq 0$ . Volume  $= \int_0^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \int_0^x dz dy dx =$   
 $\int_0^1 2x\sqrt{1-x^2} dx = [-\frac{2}{3}(1-x^2)^{3/2}]_0^1 = \frac{2}{3}$ .  
 20 Lying along the  $x$  axis the cylinder goes from  $x = 0$  to  $x = 6$ . Its slices are circular disks  $y^2 + (z-1)^2 = 1$   
 resting on the  $x$  axis. Volume  $= \int_0^6 \int_{-1}^1 \int_{1-\sqrt{1-y^2}}^{1+\sqrt{1-y^2}} dz dy dx =$  still  $6\pi$ .  
 22 Change variables to  $X = \frac{x}{a}, Y = \frac{y}{b}, Z = \frac{z}{c}$ ; then  $dX dY dZ = \frac{dx dy dz}{abc}$ . Volume  $= \iiint abc dX dY dZ =$   
 $\frac{1}{6}abc$ . Centroid  $(\bar{x}, \bar{y}, \bar{z}) = (a\bar{X}, b\bar{Y}, c\bar{Z}) = (\frac{a}{4}, \frac{b}{4}, \frac{c}{4})$ . (Recall volume  $\frac{1}{6}$  and centroid  $(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$  of standard  
 tetrahedron: this is Example 2.)  
 24 (a) Change variables to  $X = \frac{x}{4}, Y = \frac{y}{2}, Z = \frac{3z}{4}$ . Then the solid is  $X^2 + Y^2 + Z^2 = 1$ , a unit sphere of volume  
 $\frac{4\pi}{3}$ . Therefore the original volume is  $\frac{4\pi}{3}(4)(2)(\frac{4}{3}) = \frac{128\pi}{9}$ . (b) The hypervolume in 4 dimensions is  $\frac{1}{24}$ ,  
 following the pattern of 1 for interval,  $\frac{1}{2}$  for triangle,  $\frac{1}{6}$  for tetrahedron.  
 26 Average of  $f = \iiint_V f(x, y, z) dV / \iiint_V dV$  = integral of  $f(x, y, z)$  divided by the volume.  
 28 Volume of unit cube  $= \sum_{i=1}^{1/\Delta x} \sum_{j=1}^{1/\Delta x} \sum_{k=1}^{1/\Delta x} (\Delta x)^3 = 1$ .  
 30 In one variable, the midpoint rule is correct for the functions 1 and  $x$ . In three variables it is correct for  
 1,  $x, y, z, xy, xz, yz, xyz$ .  
 32 Simpson's Rule has coefficients  $\frac{1}{6}, \frac{4}{6}, \frac{1}{6}$  over a unit interval. In three dimensions the 8 corners of the cube will  
 have coefficients  $(\frac{1}{6})^3 = \frac{1}{216}$ . The center will have  $(\frac{4}{6})^3 = \frac{64}{216}$ . The centers of the 12 edges will have  
 $(\frac{1}{6})^2(\frac{4}{6}) = \frac{4}{216}$ . The centers of the 6 faces have  $(\frac{1}{6})(\frac{4}{6})^2 = \frac{16}{216}$ . (Check:  $8(1) + 64 + 12(4) + 6(16) = 216$ ).  
 When  $N^3$  cubes are stacked together, with  $N$  small cubes each way, there are only  $2N + 1$  meshpoints

along each direction. This makes  $(2N + 1)^3$  points or about 8 per cube. (Visualize the 8 new points of the cube as having  $x, y, z$  equal to zero or  $\frac{1}{2}$ .)

## 14.4 Cylindrical and Spherical Coordinates (page 547)

The three cylindrical coordinates are  $r\theta z$ . The point at  $x = y = z = 1$  has  $r = \sqrt{2}, \theta = \pi/4, z = 1$ . The volume integral is  $\iiint r dr d\theta dz$ . The solid region  $1 \leq r \leq 2, 0 \leq \theta \leq 2\pi, 0 \leq z \leq 4$  is a hollow cylinder (a pipe). Its volume is  $12\pi$ . From the  $r$  and  $\theta$  integrals the area of a ring (or washer) equals  $3\pi$ . From the  $z$  and  $\theta$  integrals the area of a shell equals  $2\pi r z$ . In  $r\theta z$  coordinates the shapes of cylinders are convenient, while boxes are not.

The three spherical coordinates are  $\rho\phi\theta$ . The point at  $x = y = z = 1$  has  $\rho = \sqrt{3}, \phi = \cos^{-1}1/\sqrt{3}, \theta = \pi/4$ . The angle  $\phi$  is measured from the  $z$  axis.  $\theta$  is measured from the  $x$  axis.  $\rho$  is the distance to the origin, where  $r$  was the distance to the  $z$  axis. If  $\rho\phi\theta$  are known then  $x = \rho \sin \phi \cos \theta, y = \rho \sin \phi \sin \theta, z = \rho \cos \phi$ . The stretching factor  $J$  is a 3 by 3 determinant and volume is  $\iiint r^2 \sin \phi dr d\phi d\theta$ .

The solid region  $1 \leq \rho \leq 2, 0 \leq \phi \leq \pi, 0 \leq \theta \leq 2\pi$  is a hollow sphere. Its volume is  $4\pi(2^3 - 1^3)/3$ . From the  $\phi$  and  $\theta$  integrals the area of a spherical shell at radius  $\rho$  equals  $4\pi\rho^2$ . Newton discovered that the outside gravitational attraction of a sphere is the same as for an equal mass located at the center.

$$\begin{array}{ll} 1 (r, \theta, z) = (D, 0, 0); (\rho, \phi, \theta) = (D, \frac{\pi}{2}, 0) & 3 (r, \theta, z) = (0, \text{any angle}, D); (\rho, \phi, \theta) = (D, 0, \text{any angle}) \\ 5 (x, y, z) = (2, -2, 2\sqrt{2}); (r, \theta, z) = (2\sqrt{2}, -\frac{\pi}{4}, 2\sqrt{2}) & 7 (x, y, z) = (0, 0, -1); (r, \theta, z) = (0, \text{any angle}, -1) \\ 9 \phi = \tan^{-1}(\frac{r}{z}) & 11 45^\circ \text{ cone in unit sphere: } \frac{2\pi}{3}(1 - \frac{1}{\sqrt{2}}) \\ 13 \text{ cone without top: } \frac{7\pi}{3} \end{array}$$

$$\begin{array}{ll} 15 \frac{1}{4} \text{ hemisphere: } \frac{\pi}{6} & 17 \frac{\pi^2}{8} \\ 23 \frac{2}{3}a^3 \tan \alpha \text{ (see 8.1.39)} & 27 \frac{\partial q}{\partial D} = \frac{\rho - D \cos \phi}{q} = \frac{\text{near side}}{\text{hypotenuse}} = \cos \alpha \end{array}$$

31 Wedges are not exactly similar; the error is higher order  $\Rightarrow$  proof is correct

$$33 \text{ Proportional to } 1 + \frac{1}{h}(\sqrt{a^2 + (D-h)^2} - \sqrt{a^2 + D^2})$$

$$35 J = \begin{vmatrix} a & & \\ & b & \\ & & c \end{vmatrix} = abc; \text{ straight edges at right angles}$$

$$37 \begin{vmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{vmatrix} = r$$

$$39 \frac{8\pi\rho^4}{3}; \frac{2}{3} \quad 41 \rho^3; \rho^2; \text{ force} = 0 \text{ inside hollow sphere}$$

$$\begin{array}{ll} 2 (r, \theta, z) = (D, \frac{3\pi}{2}, 0); (\rho, \phi, \theta) = (D, \frac{\pi}{2}, \frac{3\pi}{2}) & 4 (r, \theta, z) = (5, \cos^{-1}\frac{3}{5}, 5); (\rho, \phi, \theta) = (5\sqrt{2}, \frac{\pi}{4}, \cos^{-1}\frac{3}{5}) \\ 6 (x, y, z) = (\frac{3}{2}, \frac{\sqrt{3}}{2}, 1); (r, \theta, z) = (\sqrt{3}, \frac{\pi}{6}, 1) & 8 x = r \text{ on the positive } x \text{ axis } (x \geq 0, y = 0 (= \theta), z = 0) \end{array}$$

$$10 x = \cos t, y = \frac{\sqrt{2}}{2} \sin t, z = \frac{\sqrt{2}}{2} \sin t. \text{ The unit sphere intersects the plane } y = z.$$

$$12 \text{ The surface } z = 1 + r^2 = 1 + x^2 + y^2 \text{ is a paraboloid (parabola rotated around the } z \text{ axis). The region is above the half-disk } 0 \leq r \leq 1, 0 \leq \theta \leq \pi. \text{ The volume is } \frac{3}{4}\pi.$$

$$14 \text{ This is the volume of a half-cylinder (because of } 0 \leq \theta \leq \pi) : \text{ height } \pi, \text{ radius } \pi, \text{ volume } \frac{1}{2}\pi^4.$$

$$16 \text{ The upper surface } \rho = 2 \text{ is the top of a sphere. The lower surface } \rho = \sec \phi \text{ is the plane } z = \rho \cos \phi = 1.$$

(The angle  $\phi = \frac{\pi}{3}$  is the meeting of sphere and plane, where  $\sec \phi = 2$ .) The volume is

$$2\pi \int_0^{\pi/3} \left( \frac{8 - \sec^2 \phi}{3} \right) \sin \phi d\phi = 2\pi \left[ -\frac{8}{3} \cos \phi - \frac{1}{6 \cos^2 \phi} \right]_0^{\pi/3} = 2\pi \left[ -\frac{4}{3} - \frac{1}{6/4} + \frac{8}{3} + \frac{1}{6} \right] = \frac{5\pi}{3}.$$

- 18** The region  $1 \leq \rho \leq 3$  is a hollow sphere (spherical shell). The limits  $0 \leq \phi \leq \frac{\pi}{4}$  keep the part that lies above a  $45^\circ$  cone. The volume is  $\frac{52\pi}{3}(1 - \frac{\sqrt{2}}{2})$ .
- 20** From the unit ball  $\rho \leq 1$  keep the part above the cone  $\phi = 1$  radian and inside the wedge  $0 \leq \theta \leq 1$  radian. Volume =  $\frac{1}{4} \int_0^1 \sin \phi d\phi = \frac{1}{4}(1 - \cos 1)$ .
- 22** The curve  $\rho = 1 - \cos \phi$  is a **cardioid** in the  $xz$  plane (like  $r = 1 - \cos \theta$  in the  $xy$  plane). So we have a **cardioid of revolution**. Its volume is  $\frac{8\pi}{3}$  as in Problem 9.3.35.
- 24** Mass =  $\int_0^{2\pi} \int_0^\pi \int_0^R \rho \sin \phi (\rho + 1) d\rho d\phi d\theta = \frac{4}{3}\pi R^3 + 2\pi R^2$ .
- 26 Newton's achievement** The cosine law (see hint) gives  $\cos \alpha = \frac{D^2 + q^2 - \rho^2}{2qD}$ . Then integrate  $\frac{\cos \alpha}{q^2}$ :  $\iiint \left( \frac{D^2 - \rho^2}{2q^3 D} + \frac{1}{2qD} \right) dV$ . The second integral is  $\frac{1}{2D} \iiint \frac{dV}{q} = \frac{4\pi R^3/3}{2D^2}$ . The first integral over  $\phi$  uses the same  $u = D^2 - 2\rho D \cos \phi + \rho^2 = q^2$  as in the text:  $\int_0^\pi \frac{\sin \phi d\phi}{q^3} = \int \frac{du/2\rho D}{u^{3/2}} = [\frac{-1}{\rho D u^{1/2}}]_{\phi=0}^{\phi=\pi} = \frac{1}{\rho D} \left( \frac{1}{D-\rho} - \frac{1}{D+\rho} \right) = \frac{2}{D(D^2 - \rho^2)}$ . The  $\theta$  integral gives  $2\pi$  and then the  $\rho$  integral is  $\int_0^R 2\pi \frac{2}{D(D^2 - \rho^2)} \frac{D^2 - \rho^2}{2D} \rho^2 d\rho = \frac{4\pi R^3/3}{D^2}$ . The two integrals give  $\frac{4\pi R^3/3}{D^2}$  as Newton hoped and expected.
- 28** The small movement produces a right triangle with hypotenuse  $\Delta D$  and almost the same angle  $\alpha$ . So the new small side  $\Delta q$  is  $\Delta D \cos \alpha$ .
- 30**  $\iint q dA = 4\pi \rho^2 D + \frac{4\pi}{3} \frac{\rho^4}{D}$ . Divide by  $4\pi \rho^2$  to find  $\bar{q} = D + \frac{\rho^2}{3D}$  for the shell. Then the integral over  $\rho$  gives  $\iiint q dV = \frac{4\pi}{3} R^3 D + \frac{4\pi}{15} \frac{R^5}{D}$ . Divide by the volume  $\frac{4\pi}{3} R^3$  to find  $\bar{q} = D + \frac{R^2}{5D}$  for the solid ball.
- 32 Yes.** First concentrate the Earth to a point at its center – this is OK for each point in the Sun. Then concentrate the Sun at its center – this does not change the force on the (concentrated) Earth.
- 34**  $J = aei + bfg + cdh - ceg - afh - bdi$ .
- 36** Column 1:  $\sqrt{\sin^2 \phi(\cos^2 \theta + \sin^2 \theta)} + \cos^2 \phi = 1$ ; Column 2:  $\sqrt{\rho^2 \cos^2 \phi(\cos^2 \theta + \sin^2 \theta)} + \rho^2 \sin^2 \phi = \rho$ ; Column 3:  $\sqrt{\rho^2 \sin^2 \phi(\sin^2 \theta + \cos^2 \theta)} = \rho \sin \phi$ . These are the edge lengths of the box. The dot products of these columns are zero; so  $J = \text{volume of box} = (1)(\rho)(\rho \sin \phi)$  as before.
- 38** Column 1:  $\sqrt{\cos^2 \theta + \sin^2 \theta} = 1$ ; Column 2:  $\sqrt{r^2 \sin^2 \theta + r^2 \cos^2 \theta} = r$ ; Column 3:  $\sqrt{0^2 + 0^2 + 1^2} = 1$ . Again the dot products of the columns are zero and  $J = \text{volume of box} = (1)(r)(1) = r$ .
- 40**  $I = \frac{8}{15}\pi R^5$ ;  $J = \frac{2}{5}$ ; the mass is closer to the axis.
- 42** The ball comes to a stop at Australia and returns to its starting point. It continues to oscillate in harmonic motion  $y = R \cos(\sqrt{c/m} t)$ .

MIT OpenCourseWare  
<https://ocw.mit.edu>

Resource: Calculus  
Gilbert Strang

The following may not correspond to a particular course on MIT OpenCourseWare, but has been provided by the author as an individual learning resource.

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.